

"PRO PLUS" TOPICAL ANESTHETIC

For Use By Licensed Professionals Only

This pre-deadener was developed by a pharmacist familiar with our industries. Previous versions of this anesthetic have been used successfully for years. This recent alteration has improved its speed and depth of anesthesia. This cream contains 2% tetracaine and 4% lidocaine, has a pH of 7 and is safe for eyeliner.

External Use Only On Intact Skin. Do Not Swallow. If accidentally swallowed, get medical help immediately. Avoid contact with eyes. In case of accidental contact with eyes, rinse immediately with copious amounts of eyewash and seek care by an eye care specialist.

Keep out of reach of children. Do not use if you have a known allergy or sensitivity to any component of this product. Sensitivity test advised prior to use. If sensitivity occurs, discontinue use, immediately cleanse skin to remove product and seek medical attention as needed.

SUGGESTED APPLICATION TIMES:

Eyebrows: 15-20 minutes. First, lightly massage the cream into the brows. Then reapply a medium layer of cream over the area. Remove completely prior to procedure. Use **SJ3**, during the procedure to continue anesthesia, and either **Zone 2**, **Vasocaine** or **Blue Gel** to continue anesthesia and constrict blood vessels.

Eyeliner: 10-15 minutes. Fill the eyes with Celluvisc eye gel or other lubricant to coat the cornea. With the client's eyes closed, apply a thin layer of cream with a small applicator across the lash line. Remove anesthetic before beginning procedure. Following the first pass of tattooing, sparingly apply **Zone 2** or **Blue Gel**, to control swelling and maintain anesthesia for up to two hours.

Lips: 15-20 minutes. Lightly massage the cream into the lips. Then reapply to create a medium layer of cream over the entire area. Remove completely prior to tattooing. While some relief may be provided with topicals, for total anesthesia you must provide a nerve block performed by a physician. Do not use inside the mouth. Use **SJ3**, during the procedure to continue anesthesia, and either **Zone 2**, **Vasocaine** or **Blue Gel Anesthetic** to continue anesthesia and constrict blood vessels.

Body Skin: 60 – 90 minutes. Highly effective if left on areas above the waist for up to one hour, or below the waist for an hour and one-half prior to tattooing or piercing. Gently massage into skin and then apply a medium coating over the top. In order to prevent absorption into clothing, cover with an occlusive dressing. Thicker skin requires longer time to absorb the cream than thinner skin. Use **SJ3** or **Liquidcaine** during procedure to continue anesthesia, and either **Blue Gel**, **Vasocaine** or **Zone 2** to continue anesthesia and constrict blood vessels.

Piercing: 10-15 minutes. Gently massage into skin and then apply a medium coating on top. If the area is covered by clothing apply an occlusive dressing. Wipe away product immediately prior to piercing. Avoid genital areas. Use **Blue Gel**, **Vasocaine** or **Zone 2** during jewelry insertion to continue anesthesia and reduce swelling.